


# Turbulence in a complex plasma

Mierk Schwabe, Sergey Zhdanov, Christoph R  th

German Aerospace Center (DLR)

Knowledge for Tomorrow







# Turbulence in a complex plasma

Experiment

Signatures of Turbulence

Conclusion





# Turbulence in a complex plasma


Experiment

Signatures of Turbulence

Conclusion




We performed experiments in  
the PK-3 Plus Laboratory on the ISS.




capacitively coupled plasma chamber

We performed experiments in  
the PK-3 Plus Laboratory on the ISS.




RF voltage on electrodes

We performed experiments in  
the PK-3 Plus Laboratory on the ISS.




inject microparticles via dispensers

We performed experiments in  
the PK-3 Plus Laboratory on the ISS.



illuminate with laser from the side, trace particles

The PK-3 Plus Laboratory was hosted on the ISS from 2005 – 2014.




The PK-3 Plus Laboratory was hosted on the ISS from 2005 – 2014.



ISS014E05960

The system is often in a fluid state. Then vortices can form.






The system is often in a fluid state. Then vortices can form.




A heartbeat instability causes waves and oscillons.




A heartbeat instability causes waves and oscillons.




A heartbeat instability causes waves and oscillons.




The instability causes a drift and heating of the particles.



The particles begin to oscillate.



The flow field is quite complicated.



# Turbulence in a complex plasma

Experiment

Signatures of Turbulence

Conclusion





The Reynolds number determines the character of a fluid flow.

$$\mathcal{R} = LV/\nu$$

$L$ : characteristic length

$V$ : characteristic velocity

$\nu$ : kinematic viscosity



# The Reynolds number determines the character of a fluid flow.

$$\mathcal{R} = LV/\nu$$

$L$ : characteristic length

$V$ : characteristic velocity

$\nu$ : kinematic viscosity



# The Reynolds number determines the character of a fluid flow.

$$\mathcal{R} = LV/\nu$$

$L$ : characteristic length

$V$ : characteristic velocity

$\nu$ : kinematic viscosity

$\nu$  large,  $\mathcal{R}$  small: viscosity damps out velocity variations



# The Reynolds number determines the character of a fluid flow.

$$\mathcal{R} = LV/\nu$$

$L$ : characteristic length

$V$ : characteristic velocity

$\nu$ : kinematic viscosity

$\nu$  small,  $\mathcal{R}$  large: nonlinear effects dominate, e.g. small scale movement develops, which leads even smaller scale movement, . . .



The Reynolds number determines the character of a fluid flow.

$$\mathcal{R} = LV/\nu \approx 4$$

$L$ : characteristic length

$V$ : characteristic velocity

$\nu$ : kinematic viscosity



The Reynolds number in complex plasmas is very low.

$$\mathcal{R} = LV/\nu \approx 4$$



The Reynolds number in complex plasmas is very low.

$$\mathcal{R} = LV/\nu \approx 4$$

turbulence typically occurs at very high Reynolds numbers  $\mathcal{R}$

systems which are turbulent at low  $\mathcal{R}$  exist, e.g. viscoelastic fluids




# The Reynolds number in complex plasmas is very low.

$$\mathcal{R} = LV/\nu \approx 4$$

turbulence typically occurs at very high Reynolds numbers  $\mathcal{R}$

systems which are turbulent at low  $\mathcal{R}$  exist, e.g. viscoelastic fluids




polymer solution with  
 $\mathcal{R} = 0.7$

Groisman and Stein-  
berg, Nature (2000)






Turbulence is a superposition of movement on many scales.




NEPF Lab, MIT




We calculate velocity maps.



Care needs to be taken to make sure that holes in the velocity maps do not influence the energy spectra.




The energy spectrum is dominated by three wave numbers.



$k_{\text{fr}} = \frac{\gamma_{\text{damp}}}{2C_{\text{DAW}}} \approx 0.84 \text{ mm}^{-1}$  scale defined by friction, transition at  $k < k_{\text{fr}}$   
to  $E_k \propto k^{-3}$  caused by friction




The energy spectrum is dominated by three wave numbers.



$$k_{\text{exc}} = \frac{2\pi f_{\text{HB}}}{C_{\text{DAW}}} \approx 2.7 \text{ mm}^{-1} \text{ scale defined by heartbeat}$$




The energy spectrum is dominated by three wave numbers.



$k_{osc} = \frac{2\pi}{w_{osc}} \approx 9.0 \text{ mm}^{-1}$  scale defined by oscillons  
modulational instability causes oscillons



The knee at  $k_{\text{exc}}$  resembles a double cascade predicted for forced turbulence.




external force input at  $k_{\text{exc}}$  leads to

- inverse cascade of kinetic energy to  $k \ll k_{\text{exc}}$  ( $E \propto k^{-5/3}$ )
- direct cascade of enstrophy to  $k \gg k_{\text{exc}}$  ( $E \propto k^{-3}$ )



The knee at  $k_{\text{exc}}$  resembles a double cascade predicted for forced turbulence.




after Laurie et al. (2012)






The energy spectrum changes with the onset of the instability.



blue crosses: before / red circles: after the onset of the instability




The energy spectrum changes with the onset of the instability.



blue crosses: before / red circles: after the onset of the instability




The energy spectrum changes with the onset of the instability.



blue crosses: before / red circles: after the onset of the instability




Fitting also results in the expected power-law exponents.



linear fit:  $\log E = a - n \log k$  in both ranges separately



Fitting also results in the expected power-law exponents.



linear fit:  $\log E = a - n \log k$  in both ranges separately

We define the reduced rates of energy and enstrophy transfer.

inverse cascade:  $E = C\epsilon^{2/3}k^{-5/3} = \epsilon'^{2/3}k^{-5/3}$

$\epsilon$ : rate of cascade of kinetic energy / mass

direct enstrophy cascade:  $E = \tilde{C}\eta^{2/3}k^{-3} = \eta'^{2/3}k^{-3}$

$\eta$ : rate of cascade of mean-square vorticity (enstrophy)



We define the reduced rates of energy and enstrophy transfer.

inverse cascade:  $E = C\epsilon^{2/3}k^{-5/3} = \epsilon'^{2/3}k^{-5/3}$

$\epsilon$ : rate of cascade of kinetic energy / mass

direct enstrophy cascade:  $E = \tilde{C}\eta^{2/3}k^{-3} = \eta'^{2/3}k^{-3}$

$\eta$ : rate of cascade of mean-square vorticity (enstrophy)



We define the reduced rates of energy and enstrophy transfer.

inverse cascade:  $E = C\epsilon^{2/3}k^{-5/3} = \epsilon'^{2/3}k^{-5/3}$

$\epsilon$ : rate of cascade of kinetic energy / mass


direct enstrophy cascade:  $E = \tilde{C}\eta^{2/3}k^{-3} = \eta'^{2/3}k^{-3}$

$\eta$ : rate of cascade of mean-square vorticity (enstrophy)





The reduced rates of transfer are calculated at fixed wave numbers.



$$E = \epsilon'^{2/3} k^{-5/3}$$
$$k_1 = 2 \text{ mm}^{-1}$$

$$E = \eta'^{2/3} k^{-3}$$
$$k_2 = 6 \text{ mm}^{-1}$$



The ratio of the reduced rates of transfer indicates the excitation wave number.

$$E = \epsilon'^{2/3} k^{-5/3}$$

$$E = \eta'^{2/3} k^{-3}$$

$$\text{at } k = k_{\text{exc}}: \epsilon'^{2/3} k^{-5/3} = \eta'^{2/3} k^{-3}$$

$$k_{\text{exc}} = \sqrt{\eta'/\epsilon'} == \kappa$$




We measure  $\kappa$  as function of time.

$$\kappa^2 = \eta' / \epsilon' = \left( \frac{Ek^3|_{k_2}}{Ek^{5/3}|_{k_1}} \right)^{3/2}$$




We measure  $\kappa$  as function of time.




$$\kappa = k_{exc} = 3.9 \pm 0.5 \text{ mm}^{-1}$$




The excitation wave number determined in two separate ways agree.




$$\kappa = k_{exc} = 3.9 \pm 0.5 \text{ mm}^{-1}$$



The excitation wave number determined in two separate ways agree.



$$\kappa = k_{\text{exc}} = 3.9 \pm 0.5 \text{ mm}^{-1}$$



$$k_{\text{exc}} = \frac{2\pi f_{\text{HB}}}{C_{\text{DAW}}} \approx 2.7 \text{ mm}^{-1} \text{ scale defined by heartbeat}$$



# Turbulence in a complex plasma

Experiment

Signatures of Turbulence

Conclusion



# Conclusion

observed a double cascade in an unstable complex plasma


change in spectrum when instability sets in, fitted power-law exponents agree with those predicted for double cascade

calculated excitation wave number by two methods

still a lot to do! e.g., study fluxes, movement of particles, ...




# Conclusion




Thank you!

Knowledge for Tomorrow

# Sample Tracks



# Oscillons - frames color-coded



Enstrophy is a common quantity  
used in turbulence research.

$$\Omega_{kl} = \left( \frac{1}{n} \sum_{i=0}^n \omega_i \right)^2$$

$(k, l)$ : grid cell coordinates

$\omega_i = (\text{curl } v)_i$ : vorticity

$n$ : number of particles in cell



Enstrophy is a common quantity  
used in turbulence research.

$$\Omega_{kl} = \left( \frac{1}{n} \sum_{i=0}^n \omega_i \right)^2$$

$(k, l)$ : grid cell coordinates


$\omega_i = (\text{curl } v)_i$ : vorticity

$n$ : number of particles in cell

→ potential density related to the kinetic energy that corresponds to  
dissipation effects in the fluid



# Enstrophy map



# Enstrophy spectrum

